

# GAUGE FIELD THEORY

## Examples Sheet 1

An uppercase A, B, or C after each question (or part question) number indicates the rough degree of difficulty.

### The Klein-Gordon and Dirac equations

1. (a A) By applying the minimal substitution prescription  $p^\mu \rightarrow p^\mu - qA^\mu$  to the Klein-Gordon equation, obtain a covariant equation describing a spin zero particle of charge  $q$  in the presence of an electromagnetic potential  $A^\mu(x)$ . Show that the resulting equation is invariant under the charge conjugation operation  $\phi \rightarrow \phi^c = \phi^*$ ,  $q \rightarrow -q$ .

(b A) Apply the minimal substitution prescription to the Dirac equation, and show that the resulting equation is invariant under the charge conjugation operation  $\psi \rightarrow \psi^c = C(\bar{\psi})^T$ ,  $q \rightarrow -q$ , provided the  $4 \times 4$  matrix  $C$  satisfies the relation  $C\gamma^{\mu T}C^{-1} = -\gamma^\mu$ .

(c A) Show that, in the Pauli-Dirac representation, a suitable choice for  $C$  is  $C = i\gamma^2\gamma^0$ .

(d B) Find the result of applying the charge conjugation operation to the plane-wave solutions  $u_1(p)e^{-ip \cdot x}$  and  $u_2(p)e^{-ip \cdot x}$  defined in the lecture notes.

### Dirac gamma matrix algebra and trace theorems

2. Defining  $\not{a} \equiv \gamma^\mu a_\mu$  and using  $\gamma^\mu\gamma^\nu + \gamma^\nu\gamma^\mu = 2g^{\mu\nu}$ , prove the following results:

(a A) The trace of the product of an odd number of  $\gamma$ -matrices is zero

(b A)  $\text{Tr}(\gamma^\mu\gamma^\nu) = 4g^{\mu\nu}$ ,  $\text{Tr}(\not{a}\not{b}) = 4(a \cdot b)$

(c B)  $\text{Tr}(\gamma^\mu\gamma^\nu\gamma^\sigma\gamma^\lambda) = 4[g^{\mu\nu}g^{\sigma\lambda} + g^{\mu\lambda}g^{\nu\sigma} - g^{\mu\sigma}g^{\nu\lambda}]$   
 $\text{Tr}(\not{a}\not{b}\not{c}\not{d}) = 4[(a \cdot b)(c \cdot d) + (a \cdot d)(b \cdot c) - (a \cdot c)(b \cdot d)]$   
 $\text{Tr}(\not{a}\gamma^\mu\not{b}\gamma_\mu) = -8(a \cdot b)$

(d A)  $\gamma_\mu\not{a}\gamma^\mu = -2\not{a}$

(e A)  $\gamma_\mu\not{a}\not{b}\gamma^\mu = 4(a \cdot b)$

(f B)  $\gamma_\mu \not{a} \not{b} \not{c} \gamma^\mu = -2 \not{a} \not{b} \not{c}$ .

(g A)  $\text{Tr}(\gamma^5) = 0$

(h B)  $\text{Tr}(\gamma^5 \gamma^\mu \gamma^\nu) = 0, \quad \text{Tr}(\gamma^5 \not{a} \not{b}) = 0$

(i C)  $\text{Tr}(\gamma^5 \gamma^\mu \gamma^\nu \gamma^\lambda \gamma^\sigma) = 4i \epsilon^{\mu\nu\lambda\sigma}, \quad \text{Tr}(\gamma^5 \not{a} \not{b} \not{c} \not{d}) = 4i \epsilon^{\mu\nu\lambda\sigma} a_\mu b_\nu c_\lambda d_\sigma$

[Hint: Useful tricks are to use  $\text{Tr}(ABC) = \text{Tr}(BCA)$ , and to insert  $(\gamma^5)^2 = 1$  into a trace and then use  $\gamma^\mu \gamma^5 = -\gamma^5 \gamma^\mu$ .]

[ $\epsilon^{\mu\nu\lambda\sigma}$  is the totally antisymmetric Levi-Civita tensor defined as

$$\epsilon^{\mu\nu\lambda\sigma} = \begin{cases} -1 & \text{if } \mu\nu\lambda\sigma \text{ is an } \textit{even} \text{ permutation of } 0123, \\ +1 & \text{if } \mu\nu\lambda\sigma \text{ is an } \textit{odd} \text{ permutation of } 0123, \\ 0 & \text{if any two indices are the same.} \end{cases}$$

## Particle decay

3. The Lorentz-invariant matrix element for the decay  $\pi^- \rightarrow \mu^- \bar{\nu}_\mu$  is given by

$$M_{\text{fi}} = \frac{G_{\text{F}}}{\sqrt{2}} f_\pi p_{1\mu} \bar{u}(p_3) \gamma^\mu (1 - \gamma^5) v(p_4),$$

where  $p_1$  is the four-momentum of the  $\pi^-$ , of spin zero,  $p_3$  and  $p_4$  are the four-momenta of the  $\mu^-$  and  $\bar{\nu}_\mu$ , respectively, and  $f_\pi$  is a constant.

(a B) Show that, summed over all possible spin states of the muon, the  $\pi^- \rightarrow \mu^- \bar{\nu}_\mu$  decay rate is proportional to

$$2(p_1 \cdot p_3)(p_1 \cdot p_4) - m_\pi^2(p_3 \cdot p_4) + m_\nu [2(p_1 \cdot p_3)(p_1 \cdot s_4) - m_\pi^2(p_3 \cdot s_4)],$$

where  $s_4$  is a four-vector describing the anti-neutrino spin state, defined by

$$v(p_4) \bar{v}(p_4) \equiv (\not{p}_4 - m_\nu) \frac{1}{2} (1 + \gamma^5 \not{s}_4).$$

For a positive-helicity antineutrino, the spin four-vector  $s^\mu$  is given by

$$s_4^\mu = \frac{1}{m_\nu} (p^*, 0, 0, E_\nu^*),$$

while for a negative-helicity neutrino it has opposite sign. Hence show that the decay rate is proportional to

$$m_\mu^2(m_\pi^2 - m_\mu^2) + m_\nu^2(m_\pi^2 + 2m_\mu^2) - m_\nu^4 \pm 2(m_\mu^2 - m_\nu^2)m_\pi p^*,$$

where  $p^*$  is the momentum of the antineutrino in the  $\pi$  rest frame, and the  $+$  ( $-$ ) sign corresponds to decay into positive-helicity (negative-helicity) antineutrinos.

(b C) Assuming  $m_\nu \ll m_\pi$  and  $m_\nu \ll m_\mu$ , show that, to order  $m_\nu^2$ , the fraction of negative-helicity antineutrinos produced in  $\pi^-$  decay is given by

$$R_\downarrow \approx \frac{m_\nu^2}{m_\mu^2} \left( 1 - \frac{m_\mu^2}{m_\pi^2} \right)^{-2}.$$

Evaluate  $R_\downarrow$  for an assumed antineutrino mass of 0.1 eV.

### Quantum fields

4 A. Show that if  $\Psi$  and  $\Psi^*$  are taken as independent classical fields, the Lagrangian density

$$\mathcal{L} = \frac{\hbar}{2i} \left( \Psi \frac{\partial \Psi^*}{\partial t} - \Psi^* \frac{\partial \Psi}{\partial t} \right) - \frac{\hbar^2}{2m} \nabla \Psi \cdot \nabla \Psi^* - V(\mathbf{r}) \Psi^* \Psi$$

leads to the Schrödinger equation

$$i\hbar \frac{\partial \Psi}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \Psi + V(\mathbf{r}) \Psi$$

and its complex conjugate. What are the momentum densities conjugate to  $\Psi$  and  $\Psi^*$ ? Deduce the Hamiltonian density, and verify that integrating it over all space gives the usual expression for the energy.

### The Dirac field

5 A. The Fourier representation of the Dirac field operator is

$$\hat{\psi}(\mathbf{r}, t) = \int \frac{d^3 \mathbf{k}}{(2\pi)^3 2\omega} \sum_s \left[ \hat{c}_s(\mathbf{k}) u_s(\mathbf{k}) e^{-ik \cdot x} + \hat{d}_s^\dagger(\mathbf{k}) v_s(\mathbf{k}) e^{+ik \cdot x} \right]$$

where  $k^\mu = (\omega, \mathbf{k})$  and  $\omega = \sqrt{\mathbf{k}^2 + m^2}$ . The creation and annihilation operators satisfy the anticommutation relations

$$\{\hat{c}_s(\mathbf{k}), \hat{c}_{s'}^\dagger(\mathbf{k}')\} = \{\hat{d}_s(\mathbf{k}), \hat{d}_{s'}^\dagger(\mathbf{k}')\} = (2\pi)^3 2\omega \delta_{ss'} \delta^3(\mathbf{k} - \mathbf{k}'),$$

all other anticommutators being zero. Show that this implies that the field and its conjugate momentum density satisfy the anticommutation relation

$$\{\hat{\psi}_\alpha(\mathbf{r}, t), \hat{\pi}_\beta(\mathbf{r}', t)\} = i \delta_{\alpha\beta} \delta^3(\mathbf{r} - \mathbf{r}')$$

where  $\hat{\pi} = i\hat{\psi}^\dagger$ , and  $\alpha$  and  $\beta$  are spinor component labels.

Show also that the normal-ordering properties of the creation and annihilation operators lead to the relation

$$:\hat{\psi}_\beta(y) \hat{\psi}_\alpha(x): = -:\hat{\psi}_\alpha(x) \hat{\psi}_\beta(y):$$

for the normal-ordered products of field operator components at spacetime points  $x$  and  $y$ .

### The charge operator for a Dirac field

- 6 B. Show that the standard particle and antiparticle spinors  $u_s(\mathbf{k})$ ,  $v_s(\mathbf{k})$  in the Pauli-Dirac representation satisfy the relations

$$u_s^\dagger(\mathbf{k})u_{s'}(\mathbf{k}) = v_s^\dagger(\mathbf{k})v_{s'}(\mathbf{k}) = 2\omega\delta_{ss'}$$

$$u_s^\dagger(\mathbf{k})v_{s'}(-\mathbf{k}) = v_s^\dagger(\mathbf{k})u_{s'}(-\mathbf{k}) = 0$$

For a free spin-half particle described by a field operator  $\hat{\psi}(x)$ , verify that the current

$$\hat{J}^\mu(x) = \hat{\bar{\psi}}(x)\gamma^\mu\hat{\psi}(x)$$

is conserved, and show that it leads to a conserved (normal-ordered) charge operator

$$:\hat{Q}: = \int d^3\mathbf{k} N(\mathbf{k}) \sum_s \left[ \hat{c}_s^\dagger(\mathbf{k})\hat{c}_s(\mathbf{k}) - \hat{d}_s^\dagger(\mathbf{k})\hat{d}_s(\mathbf{k}) \right] .$$

Show that the single-particle states  $\hat{c}_s^\dagger(\mathbf{k})|0\rangle$  and  $\hat{d}_s^\dagger(\mathbf{k})|0\rangle$  are eigenstates of  $:\hat{Q}:$  and obtain the corresponding eigenvalues.